

FOR FLU AND COLD

COMPOSITION:

Syrup:

Each 5 ml contains Triprolidine Hydrochloride (BP) 1.25 mg and Pseudoephedrine Hydrochloride (BP) 30 mg.

Other ingredients: Glycerol, Sorbitol Solution, Saccharin Sodium, Sodium Methyl Hydroxybenzoate, Sodium Benzoate, Ethanol, Sodium Chloride, Cherry Flavour and Purified Water.

CLINICAL PHARMACOLOGY:

FLU-CARE treats flu symptoms:

It liberates rapidly the nose in approximately 30 minutes for the decongestant action. It decreases sneezing for its anti-allergic action.

INDICATIONS:

FLU-CARE could be used for the symptomatic treatment of acute or allergic rhinitis after exposition to flower pollens or animal hair, house dust, flu, head cold rhinopharyngitis and flu states with nose running.

CONTRAINDICATIONS:

Triprolidine Hydrochloride and Pseudoephedrine Hydrochloride should not be used:

- In patients under M.A. O.I. treatments (anti-depressant) or furazolidone treatment (urinary antiseptic).
- In patients suffering from serious hepatic disorders,
- glaucoma with angle closure, hypersensitivity to one of the components and in case of prostate adenoma.
- Treatment of cough and cold symptoms in children under 6 years of age.

PRECAUTIONS:

- Should consult the physician before using in cases of: Cardiac disease, hyperthyroidism, diabetes, psychosis, hypertension and renal or hepatic disorders.
- Triprolidine Hydrochloride and Pseudoephedrine Hydrochloride should not be used by patients taking sleeping pills (specially barbiturates) or taking alcohol.

WARNING:

- If symptoms do not improve within 7 days or are accompanied by fever, consult the physician.
- Do not dispense to children less than six years of age.

ADVERSE REACTIONS:

This medicine may cause few side effects as:

Cutaneous reactions which disappear after treatment interruption, palpitation, dry mouth, nausea, vomiting, anxiety, sleeping problems and excitations which disappear after treatment interruption.

USE DURING PREGNANCY AND LACTATION:

Should consult your doctor before use during pregnancy and lactation.

DOSAGE:

- **Adults and children over 12 years:** 10 ml of syrup 3 times a day.
- **Children from 6 to 12 years:** 5 ml of syrup 3 times a day.

Do not dispense to children less than six years of age.

DURATION OF TREATMENT:

- The average duration is 5 to 6 days.
- Treatment could be interrupted at symptoms disappearance.

STORAGE:

Store at room temperature (15-25)°C.

Do not use the drug after the expiry date printed on the package.

PRESENTATION:

Syrup: 100 ml bottles.

To Report any side effect(s) or Quality Problems:

- National Pharmacovigilance Center (NPC)
- Fax: +966-1-205-7662
- E-mail: npc.drug@sfd.a.gov.sa
- website: www.sfd.a.gov.sa/npc

This is a medicament

- A medicament is a product, which affects your health, and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are experts in medicine, its benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.

Keep medicaments out of the reach of children

Council of Arab Health Ministers
Union of Arab Pharmacists



Manufactured by **RIYADH PHARMA**
Medical and Cosmetic Products Co. Ltd.
Riyadh, Saudi Arabia.