



Buscopan

Hyoscine - N - butylbromide



Properties :

Has a special site of action at the parasympathetic ganglia in the walls of the viscera. Because of this it exerts a specific antispasmodic action on the smooth muscle of the gastro-intestinal, biliary and urinary tracts. No side effects on the central nervous system, the eye, the salivary glands or heart have been observed when Buscopan is administered in therapeutic doses.

Indications :

Gastro intestinal spasm and hypermotility, pylorospasm in infants, post-operative vomiting, spastic constipation, biliary dyskinesia and spasm of the urinary tract. Delayed relaxation of the lower uterine segment, dysmenorrhoea.

Dosage :

Adults and school children : The usual dose is 1-2 sugar coated tablets or 1-2 suppositories (each containing 10 mg) administered 3-5 times daily.

The dose can be raised at the discretion of the physician. The tablets should be swallowed whole with a little water.

The suppositories should be unwrapped and inserted into the rectum pointed end first.

In more pronounced conditions of pain 1 ampoule containing 20 mg may be injected subcutaneously, intramuscularly or intravenously several times daily.

In severe biliary and renal colic the slow intravenous injection of Buscopan compositum rapidly relieves the severe pain.

Infants and young children :

1 paediatric suppository 3 times daily.

In severe case $\frac{1}{4}$ of an ampoule 3 times daily.

Pack :

Sugar coated tablets : Each containing 10 mg	Box of 20 tablets
Suppositories for adult containing 10 mg	Box of 6 suppositories
Paediatric suppositories containing 7.5 mg	Box of 6 paediatric suppositories
Ampoules containing 20 mg	Box of 6 ampoules

Storage :

Keep at room temperature (15 - 30)°C, in cool and dry place.

Product of : **CHEMICAL INDUSTRIES DEVELOPMENT (CID)**
GIZA - A.R.E. - G.C.R. 19717

Under licence of : **Boehringer Ingelheim International GmbH**
Ingelheim am Rhein - Germany

B 203050 B
7 / 09